


Claims

1. A compound of formula (1):



wherein:

Z is CH or nitrogen;

R⁴ and R⁵ together are either -S-C(R⁶)=C(R⁷)- or -C(R⁷)=C(R⁶)-S-;

R⁶ and R⁷ are independently selected from hydrogen, halo, nitro, cyano, hydroxy,

10 fluoromethyl, difluoromethyl, trifluoromethyl, trifluoromethoxy, carboxy, carbamoyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy and (1-4C)alkanoyl;

A is phenylene or heteroarylene;


n is 0, 1 or 2;

R¹ is independently selected from halo, nitro, cyano, hydroxy, carboxy, carbamoyl,

15 N-(1-4C)alkylcarbamoyl, N,N-((1-4C)alkyl)₂carbamoyl, sulphamoyl, N-(1-4C)alkylsulphamoyl, N,N-((1-4C)alkyl)₂sulphamoyl, -S(O)_b(1-4C)alkyl (wherein b is 0,1, or 2), -OS(O)₂(1-4C)alkyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (1-4C)alkanoyl, (1-4C)alkanoyloxy, hydroxy(1-4C)alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, trifluoromethoxy and -NHSO₂(1-4C)alkyl;

20 or, when n is 2, the two R¹ groups, together with the carbon atoms of A to which they are attached, may form a 4 to 7 membered saturated ring, optionally containing 1 or 2 heteroatoms independently selected from O, S and N, and optionally being substituted by one or two methyl groups;

r is 1 or 2; and when r is 1 the group



25

is a substituent on carbon (2) and when r is 2 (hereby forming a six membered ring) the same group is a substituent on carbon (2) or on carbon (3);

- Y is selected from $-C(O)R^2$, $-C(O)OR^2$, $-C(O)NR^2R^3$, $-(1-4C)alkyl$ [optionally substituted by 1 or 2 substituents independently selected from hydroxy, $-C=NR^2$, $(1-4C)alkoxy$, aryloxy, heterocycloxy, $-S(O)_bR^2$ (wherein b is 0, 1 or 2), $-O-S(O)_bR^2$ (wherein b is 0, 1 or 2), $-NR^2R^3$, $-N(OH)R^2$, $-NR^2C(=O)R^2$, $-NHOHC(=O)R^2$, $-SO_2NR^2R^3$, $-N(R^2)SO_2R^2$, aryl and heterocyclyl], $-C(O)NOH$, $-C(O)NSH$, $-C(N)OH$, $-C(N)SH$, $-SO_2H$, $-SO_3H$, $-SO_2N(OH)R^2$, $-(2-4C)alkenyl$, $-SO_2NR^2R^3$, $-(1-4C)alkylC(O)R^2$, $-(1-4C)alkylC(O)OR^2$, $-(1-4C)alkylSC(O)R^2$, $-(1-4C)alkylOC(O)R^2$, $-(1-4C)alkylC(O)NR^2R^3$, $-(1-4C)alkylOC(O)OR^2$, $-(1-4C)alkylN(R^2)C(O)OR^2$, $-(1-4C)alkylN(R^2)C(O)NR^2R^3$, $-(1-4C)alkylOC(O)NR^2R^3$, $(3-6C)cycloalkyl$ (optionally substituted by 1 or 2 R^8), aryl, heterocyclyl (wherein the heterocyclic ring is linked by a ring carbon atom), $-(1-4C)alkylSO_2(2-4C)alkenyl$ and $-S(O)_cR^2$ (wherein c is 0, 1 or 2); R^2 and R^3 are independently selected from hydrogen, $-O(1-4C)alkyl$, $-S(1-4C)alkyl$, $-N(1-4C)alkyl$, heterocyclyl, aryl, and $(1-4C)alkyl$ [optionally substituted by 1 or 2 R^8 groups]; or
- wherein NR^2R^3 may form a 4 to 7 membered saturated, partially saturated or unsaturated ring, optionally containing 1, 2 or 3 additional heteroatoms independently selected from N, O and S (provided there are no O-O, O-S or S-S bonds), wherein any $-CH_2-$ may optionally be replaced by $-C(=O)-$, and any N or S atom may optionally be oxidised to form an N-oxide or SO or SO_2 group respectively, and wherein the ring is optionally substituted by 1 or 2 substituents independently selected from halo, cyano, $(1-4C)alkyl$, hydroxy, $(1-4C)alkoxy$ and $(1-4C)alkylS(O)_b-$ (wherein b is 0, 1 or 2); R^8 is independently selected from hydrogen, hydroxy, $(1-4C)alkyl$, $(2-4C)alkenyl$, $(1-4C)alkoxy$, cyano((1-4C))alkyl, amino((1-4C))alkyl [optionally substituted on nitrogen by 1 or 2 groups selected from $(1-4C)alkyl$, hydroxy, hydroxy((1-4C))alkyl, dihydroxy((1-4C))alkyl, $-CO_2(1-4C)alkyl$, aryl and aryl((1-4C))alkyl], halo((1-4C))alkyl, dihalo((1-4C))alkyl, trihalo((1-4C))alkyl, hydroxy((1-4C))alkyl, dihydroxy((1-4C))alkyl, $(1-4C)alkoxy(1-4C)alkoxy$, $(1-4C)alkoxy(1-4C)alkyl$, hydroxy(1-4C)alkoxy, 5- and 6-membered cyclic acetals and mono- and di-methyl derivatives thereof, aryl, heterocyclyl, (heterocyclyl)(1-4C)alkyl, $(3-7C)cycloalkyl$ (optionally substituted with 1 or 2 hydroxy groups, $(1-4C)alkyl$ or $-CO_2(1-4C)alkyl$), $(1-4C)alkanoyl$, $(1-4C)alkylS(O)_b-$ (wherein b is 0, 1 or 2), $(3-6C)cycloalkylS(O)_b-$ (wherein b is 0, 1 or 2), arylS(O)_b- (wherein b is 0, 1 or 2), heterocyclylS(O)_b- (wherein b is 0, 1 or 2), benzylS(O)_b- (wherein b is 0, 1 or 2),

(1-4C)alkylS(O)_c(1-4C)alkyl- (wherein c is 0, 1 or 2), -N(OH)CHO, -C(=N-OH)NH₂,
 -C(=N-OH)NH(1-4C)alkyl, -C(=N-OH)N((1-4C)alkyl)₂, -C(=N-OH)NH(3-6C)cycloalkyl,
 -C(=N-OH)N((3-6C)cycloalkyl)₂, -COCOOR⁹, -C(O)N(R⁹)(R¹⁰), -NHC(O)R⁹,
 -C(O)NHSO₂((1-4C)alkyl), -NHSO₂R⁹, (R⁹)(R¹⁰)NSO₂-, -COCH₂OR¹¹, -COCH₂OH,
 5 (R⁹)(R¹⁰)N-, -COOR⁹, -CH₂OR⁹, -CH₂COOR⁹, -CH₂OCOR⁹, -CH₂CH(CO₂R⁹)OH, -
 CH₂C(O)NR⁹R¹⁰, -(CH₂)_wCH(NR⁹R¹⁰)CO₂R^{9'} (wherein w is 1, 2 or 3), and
 -(CH₂)_wCH(NR⁹R¹⁰)CO(NR^{9'}R^{10'}) (wherein w is 1, 2 or 3);

R⁹, R^{9'}, R¹⁰ and R^{10'} are independently selected from hydrogen, hydroxy, (1-4C)alkyl
 (optionally substituted by 1 or 2 R¹¹), (2-4C)alkenyl, (3-7C)cycloalkyl (optionally substituted
 10 by 1 or 2 hydroxy groups), cyano((1-4C))alkyl, trihaloalkyl, aryl, heterocyclyl,
 heterocyclyl((1-4C)alkyl), -CO₂(1-4C)alkyl; or

R⁹ and R¹⁰ together with the nitrogen to which they are attached, and/or R^{9'} and R^{10'} together
 with the nitrogen to which they are attached, form a 4- to 6-membered ring where the ring is
 optionally substituted on carbon by 1 or 2 substituents independently selected from oxo,
 15 hydroxy, carboxy, halo, nitro, cyano, carbonyl, (1-4C)alkoxy and heterocyclyl; or the ring may
 be optionally substituted on two adjacent carbons by -O-CH₂-O- to form a cyclic acetal
 wherein one or both of the hydrogens of the -O-CH₂-O- group may be replaced by a methyl;
 R¹¹ is independently selected from (1-4C)alkyl and hydroxy(1-4C)alkyl;
 or a pharmaceutically acceptable salt or pro-drug thereof.

20

2. A compound of the formula (1), or a pharmaceutically acceptable salt or pro-drug
 thereof, as claimed in claim 1, wherein A is phenylene.

3. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo
 25 hydrolysable ester thereof, as claimed in claim 1 or claim 2, wherein n is 0.

4. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo
 hydrolysable ester thereof, as claimed in any one of the preceding claims wherein r is 1.

30 5. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo
 hydrolysable ester thereof, as claimed in any one of the preceding claims wherein R⁶ and R⁷
 are independently hydrogen or halo.

6. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in any one of the preceding claims wherein Y is selected from $-C(O)OR^2$, $-C(O)NR^2R^3$, $-(1-4C)alkyl$ [optionally substituted by a substituent selected from hydroxy, $(1-4C)alkoxy$, $-S(O)_bR^2$ (wherein b is 0, 1 or 2), $-O-S(O)_bR^2$ (wherein b is 0, 1 or 2), $-NR^2R^3$, $-NR^2C(=O)R^2$ and $-SO_2NR^2R^3$], $-(1-4C)alkylC(O)R^2$, $-(1-4C)alkylC(O)OR^2$, $-(1-4C)alkylOC(O)R^2$, $-(1-4C)alkylC(O)NR^2R^3$, $-(1-4C)alkylOC(O)OR^2$, $-(1-4C)alkylN(R^2)C(O)OR^2$, $-(1-4C)alkylN(R^2)C(O)NR^2R^3$, $-(1-4C)alkylSC(O)R^2$, $-(1-4C)alkylOC(O)NR^2R^3$, $-(1-4C)alkylSO_2(2-4C)alkenyl$ and $-SO_cR^2$ (wherein c is 0, 1 or 2).

10

7. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in any one of the preceding claims wherein R^2 and R^3 are independently selected from hydrogen, heterocyclyl, $-O(1-4C)alkyl$, $-N(1-4C)alkyl$, $(1-4C)alkyl$ [optionally substituted by 1 or 2 R^8 groups]; or an NR^2R^3 group forms a morpholine, thiomorpholine (and oxidised versions thereof), pyrrolidine, or piperidine ring and wherein the ring is optionally substituted by 1 or 2 substituents independently selected from chloro, fluoro, hydroxy and methoxy.

15

8. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in any one of the preceding claims wherein R^8 is independently selected from hydrogen, hydroxy, $-C(O)N(R^9)(R^{10})$, $-NHC(O)R^9$, $-COOR^9$, $-CH_2OR^9$, $-CH_2COOR^9$, $-CH_2OCOR^9$, aryl, heterocyclyl, and 5- and 6-membered cyclic acetals and mono- and di-methyl derivatives thereof.

20

9. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in any one of the preceding claims wherein R^9 and R^{10} are independently selected from hydrogen, hydroxy and $(1-4C)alkyl$ or R^9 and R^{10} together with the nitrogen to which they are attached form a morpholine, thiomorpholine (and oxidised versions thereof), pyrrolidine, or piperidine ring.

25

10. A pharmaceutical composition which comprises a compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in claim 1 in association with a pharmaceutically-acceptable diluent or carrier.

30

11. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in claim 1, for use in a method of treatment of a warm-blooded animal such as man by therapy.

5 12. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in claim 1, for use as a medicament.


13. A compound of the formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in claim 1, for use as a medicament in the treatment of
10 type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac ischaemia or obesity in a warm-blooded animal such as man.

14. The use of a compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in claim 1, in the manufacture of a medicament for
15 use in the treatment of type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac ischaemia or obesity in a warm-blooded animal such as man.

15. The use of a compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in claim 1, in the manufacture of a medicament for
20 use in the treatment of type 2 diabetes in a warm-blooded animal such as man.

16. A process for the preparation of a compound of formula (1) as claimed in claim 1, which process comprises:


reacting an acid of the formula (2):



(2)

or an activated derivative thereof; with an amine of formula (3):

- 115 -



(3)

and thereafter if necessary:

- i) converting a compound of the formula (1) into another compound of the formula (1);
- 5 ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.